

Study and Evaluation of Crude Drugs

Dr. Satyender Kumar

Associate professor Chemistry, Govt college for Women, Hisar

Email skyadavqchisar@gmail.com

Abstract

Evaluation is the methodical assessment of a subject's quality, value, and relevance in accordance with predetermined criteria and norms. Aims, realisable concepts/proposals, and alternatives may be evaluated; the feasibility of an action or intervention can be determined; and decision-making can be aided. The extent to which the goals, objectives, and outcomes of the activity have been met or exceeded. As important as it is to obtain understanding of current or past efforts, evaluation's main goal is to promote self-reflection and help pinpoint areas in need of improvement.

Keywords: crude drugs, evaluation of crude drugs, organoleptic evaluation etc.

Introduction

Several other definitions have been proposed for the fields of ethno-botany and ethno-pharmacology. It's crucial for expanding our understanding of the plants revered in tribal mythology, the variety of species they've gathered for nutrition, and the methods they've developed for harvesting and preparing them. Traditional methods of illness detection and care have evolved throughout the tribes. New pharmaceutical medications derived mostly from medicinal plants. Evaluation of a crude medicine entails establishing its authenticity and measuring its potency and purity. Neelakurinji is the Malayalam name for the plant *Chassalia curviflora* (Wall. ex Kurz.) Thwaites. Among the biggest blooming plants, it is from the family Rubiaceae. A shrub that may grow up to two metres in height, is evergreen, with elliptic ovate leaves, flowers that are pinkish white with a yellow throat, and drupes that are a deep purple [1]. The Western Ghats of Kerala, India, employ it as a traditional medicine. The Aarukani Hills' Kani natives traditionally employed *C. curviflora*'s root and root bark to cure jaundice and heal wounds[2]. Moreover, it has been stated that the plant's many components may be utilised to treat a wide range of medical conditions, including but not limited to: headaches, ulcers, sore throats, phlegm, rheumatism, pneumonia, eye illnesses, and ear infections [3, 4]. There is a dearth of information on the quantification of this plant's physicochemical and biochemical properties at the current time. Thus, this research aimed to evaluate quality control criteria for future pharmacognostical standardisation of this plant material. The only processing that has been done to the ingredients in crude medications is collecting them and drying them out. Natural substances are those that occur in nature without undergoing any alterations to their molecular structure by humans. People take use of their medicinal properties. Both human and animal patients by using the Senna or Cinchona leaf ingested or topically to treat a variety of ailments.

To diagnose, treat, mitigate, or prevent illness in humans or other animals, a crude medication is any naturally occurring, unrefined chemical produced from organic or inorganic sources such plants,

animals, microorganisms, organs, or complete organisms. [1] Color, smell, taste, size, form, and unique characteristics like texture are all taken into account throughout the organoleptic and morphological assessment process.

Identifying the pure powdered medication requires a microscope. More information about a subject may be gleaned using this approach, medication and identifying them based on their histological characteristics. The microscope's magnifying power makes it possible to expand tiny research samples, which is essential for calculating parameters like leaf constants, stomatal index, and palisade ratio. Biological: Bioassays measure medication pharmacological activity. The biological impact of the medicine is the effect it has on living creatures, such as bacteria, fungal growth, or animal tissue, that is used to determine its potency relative to the standard drug. These techniques allow for the evaluation of an unprocessed medication and the recommendation of next steps in clinical testing.

To evaluate efficacy and identify active ingredients, chemical tests perform the best. There are several analyses and examinations included. Methods of analysis include the separation, purification, and determination of bioactive components.

The approaches include quantitative chemical tests like the acid value, saponification value, etc. Active principles are modified by use of physical constants. Moisture content, specific gravity, density, optic rotation, etc., are all evaluated more accurately using these. [2]

CHEMICAL EVALUATION

The chemical evaluation includes not only quantitative and qualitative analyses of chemicals, but also chemical tests and instrumental analysis. Separation, purification, and identification of valuable components are all part of the chemical procedures used in analysis. Species identification assays and testing for phytoconstituents, such as alkaloids, glycosides, tannins, and others, are some of the qualitative chemical tests that may be performed. The procedures for determining the various phytoconstituents are explained in the various chapters of the book, and references are supplied to those chapters. Copper acetate is a useful tool for identifying colophony, an adulterant that may be discovered in resins, balsams, and waxes. In a similar manner, cottonseed oil may be recognised using the Holphen's test, and sesame oil can be identified using the Baudouin's test when it is mixed with olive oil. It is possible to distinguish gurjun balsam in copaiba by using acetic and nitric acids.

In order to evaluate the effectiveness of a treatment via the use of chemical processing, there are a few different quantitative chemical tests that may be carried out. They include the acid value (present in resins and balsams), the saponification value (present in balsams), the ester value (present in balsams and volatile oils), and the acetyl value (present in volatile oils, among other things).

The tests for alkaloids, resins, volatile oils, glycosides, vitamins, and other components are all examples of different sorts of chemical analysis. The belladonna plant, ipecacuanha, nux vomica, jalap, cod-liver oil, and strychnine are all examples of substances that have been analysed to determine whether or not they contain alkaloids, phenolics, vitamins, or any other kind of chemicals. The findings may lead one to the conclusion that the drug in question is of poor quality or has been depleted, and if the substance that was tested is not there, this may suggest that a replacement that is wholly useless has been employed.

In order to investigate the different chemical classes of phytoconstituents, several instrumental studies, such as chromatographic and spectroscopic procedures, are used. Chromatography may

be used in a variety of different ways to separate chemicals. There are several different kinds of spectroscopy, including nuclear magnetic spectroscopy, spectroscopy of the visible and ultraviolet spectrum, infrared spectroscopy, and mass spectroscopy. [3]

PHYSICAL EVALUATION

Physical approaches are often used in crude plant evaluation. These techniques may be used to measure the solubility, relative density, optical rotational, viscous, index of refraction, heat resistance, moisture content, amount of fibre elasticity, and other physical properties of the herb material.

Solubility

Drugs' individual responses to solvents are taken into consideration. This is useful for analysing several oil and oleoresin samples. Colophony can be dissolved in light petroleum, balsam of Peru can be dissolved in chloral hydrate solution, castor oil can be dissolved in half its volume of light petroleum, and turbidity can be produced by using two volumes of the solvent, but only three volumes of 90% alcohol, and so on. It's possible that alkaloidal bases may be dissolved in organic solvents, whereas alkaloidal salts can be dissolved in polar solvents.

The Optical Twist

Anisotropic crystalline materials and samples containing an excess of one homologue of a chiral compound may distort the orientation of plane-polarized light. Certain chemicals are optically active, and one of the characteristics they display is rotation in the optical spectrum. In the direction of incident light, dextrorotatory (d) or (+) enantiomers cause the light to rotate clockwise, whereas levorotatory (l) or (–) enantiomers cause the light to spin anticlockwise. For example, you may use eucalyptus oil between 0 and 10 degrees Celsius, honey between 3 and 15 degrees Celsius, Che- nopodium oil between 30 and 80 degrees Celsius, etc.

Measurement of the Refractive Index

The refractive index is the ratio of the movement of light in a vacuum to the light's speed in the substance. When a light stream is incident on one material and transmitted through the other at an angle, the refraction and transmission angles are dependent on the refraction of the two materials. As the refractive index changes with the velocity of the light, the speed at which different colours travel might vary. The refractive index may also be influenced by the intensity of the incident light. An appropriate measure for determining the efficacy of herbal medications might be their particular values, such as the range from 1.4758 to 1.527 for castor oil, 1.527 to 1.535 for clove oil, etc.

Gravity's Finer Points

Density in relation to another substance is another name for it. The comparison of the mass of a solid, liquid, or gas to that of the same volume of distilled water at 4 degrees Celsius (39 degrees Fahrenheit), or of air or hydrogen at the same temperature and pressure. Cottonseed oil ranges in specific gravity from 0.88 to 0.93, whereas coconut oil is 0.925 and castor oil is 0.95.

BIOLOGICAL EVALUATION

The pharmacological activity, potency, and toxicity of the plant or extract may then be assessed using a number of different biological approaches. Medications that could not be adequately tested by physical and chemical procedures would benefit more from a

biological review. Moreover, this is an key assessment tool, the crude medications are only deemed relevant because of their biological impacts, and this analysis would conclude the effect. It is generally agreed that these approaches are less reliable, take longer, and cost more. The screening of a wide range of biological features requires bioassays to be as straightforward as feasible, and researchers should make every effort to get access to a broad panel of tests. There are three distinct kinds of bioassays, including those that are toxicological, symptomatic, and tissue/organ based. Three distinct approaches are used: the toxicological, the symptomatic, and the isolated organ or tissue approaches.

The effective dose of a medicine is established by measuring its impact on appropriate test animals or organs under controlled laboratory circumstances. Certain types of bioassays use the use of reference standards in an effort to reduce the potential for mistakes.

To determine both the fatal and effective doses of crude drugs, toxicity studies are conducted in appropriate animal models. It is common practise to evaluate potential vaccinations on mice.

Both guinea pigs and young domestic chickens may have their oxytocic activity of vasopressin examined by injecting it into an exposed crural or brachial vein and watching the resulting changes in blood pressure. One way to assess the effectiveness of Digitalis glycosides is to inject the drug into a pigeon's alar vein and observe the bird until it dies. Various medications being tested to see whether they might calm cats and reduce mydriasis in their eyes. Anti-helminthic drugs are evaluated in worms. Rabbits' eyes are used as test subjects for eye medications. Medications with cardiac and gastrointestinal actions are tested in dogs. The uterus or intestine of a rabbit or a cock's comb are used to study Ergot's effects. Experiments are performed not just on animals, but also on human volunteers. It's possible that human research might provide results different from those shown in animal studies. Hepatoprotective, hypoglycemic, antiinflammatory, antiulcer, immunomodulatory, and other biological actions have all been studied.

Antimicrobial medications are identified via the use of microbiological tests, which examine the effects of antibiotics on a variety of microbes. Agar well-diffusion, disc-diffusion, and turbidimetric techniques are all used in these sorts of experiments. When it comes to vitamin assays, yeast moulds, which are alive and comprised of bacteria, are the technique of choice.

Cultivation and Collection

Often reaching heights of up to 6 metres in height, acacia is a very prickly tree. Gum is harvested from wild plants in Senegambia, whereas in Sudan it is extracted from specially planted trees due to the region's severe environment. Planting acacia trees is as simple as scattering the seeds over the depleted ground. It's not only the wind that spreads the trees' seeds; they may do it naturally, too. Two times a year, in dry conditions,

in November and February–March, locals pick gum from trees that are six to eight years old. Axe cuts 2-3 feet long and 2-3 inches wide are made in the stem and branches, the bark is loosened and removed by axe, and care is taken to avoid damaging the cambium and xylem. The xylem is often covered in a thin coat of bark after they are done. White ants may invade a plant if the xylem is exposed, preventing the formation of gum. In the winter, gum oozes out 6-8 weeks after damage, but in the summer it takes just 3-4 weeks. It's thought that in the warmer months, bacteria that make it through the incision are more active, leading to a faster rate of gum production. The emitted gum is cleaned by scraping off bark and wood debris, sifting off sand, etc., and then storing it in leather bags. For roughly three weeks, gum is sun-dried in trays in thin layers, which allows for the bleaching process that turns the gum whiter. This causes the gum to compress unevenly, leading to the appearance of fractures and fissures on its exterior and ultimately the loss of its translucence. The gum ripens during this process.

Conclusion

Adolescent drug abuse and criminal behaviour are closely intertwined, making identification of drug-using youth in the juvenile justice system an essential first step in acting on both issues. Drug detection methods, when applied to the existing high rates of drug usage and criminal behaviour, may help lower their prevalence. Drug testing may be a helpful intervention for teens because it forces them to face the reality of their substance abuse problem head-on. In order to help young people avoid drug use and potentially harmful actions, drug testing might be included into an intervention. Juvenile justice agencies may better fulfil their missions of community safety, teen accountability, and skill development with the help of long-term, precise drug detection.

This Overview describes the drug detection procedures that have been adopted in five facilities whose major emphasis is the supervision of adolescents on probation, as well as three juvenile detention facilities. The ACA/IBH and the APPA have provided assistance to these jurisdictions as they developed drug testing and intervention programmes that meet or exceed federal requirements. Results from teenage drug testing showed wide variation, ranging from 10% at one demonstration location to 37% at another, which is consistent with data showing widespread drug use among juvenile delinquency system juveniles. At every place we checked, marijuana was by far the most prevalent drug we found. The rate of positive results for cocaine was second highest overall. In all but one of the locations, the favourable outcomes associated with cocaine were far lower than those associated with marijuana. There were many positive PCP tests in two different places.

References

- [1]. Cooper TH, Clark G, Guzinski J (1994) Teas, spices and herbs. In: Food Phytochemicals, ed. Ho, CT, Washington, DC: American Chemical Society 1: 231 - 236.
- [2]. Allen PC, Danforth HD, Augustine PC (1998) Dietary modulation of avian coccidiosis. *Int J Parasitol* 28(7): 1131-1140.
- [3]. Mazumder A, Wang S, Neamati N, Nicklaus M, Sunder S et al. (1996)
- [4]. Antiretroviral agents as inhibitors of both human immunodeficiency virus type 1 integrase and protease. *J Med Chem* 39(13): 2472 - 2481.
- [5]. Ammon HPT, Wahl MA (1991) Pharmacology of *Curcuma longa*. *Planta Med* 57(1): 1-7
- [6]. Chauhan SK, Singh BD, Agrawala S (1999) Estimation of curcuminoids in *Curcuma longa* by HPLC and spectrophotometric methods. *Indian J Pharm Sci* 61(1): 58-60.
- [7]. Dwivedi AK, Raman M, Seth RK, Sarin JP (1993) Combined thin-layer chromatography- densitometry for the quantitation of curcumin in pharmaceutical dosage forms and in serum. *Indian J Pharm Sci* 55(5): [8]. 174-177.
- [8]. Pundarikakshudu K, Dave HN (2010) Simultaneous Determination of curcumin and berberine in their pure form and from the combined extracts of *Curcuma longa* and *Berberis aristata*. *Int J Appl Sci*
- [9]. Sahani S, Mall TP- Diversity of potent ethno-medicinal flora for headache from north-Tarai forest of (U.P.) India. *Inter. J Interdis and Multidisci Stud* 2013; 1:33-46
- [10] Pathak NN, Kamra DN, Agarwal N. Analytical techniques in animal nutrition research, Istedn. IBD. Co. 1996; 5-23
- [11] James CS. Analytical chemistry of food Seale- Hayne Faculty of Agriculture, Food and Land use, Department of Agriculture and Food studies, University of Plymouth, UK, 1995, 96-97